

Is the Mantel correlogram powerful enough to be useful in ecological  
analysis? A simulation study

DANIEL BORCARD<sup>1,2</sup> AND PIERRE LEGENDRE<sup>1</sup>

<sup>1</sup> Département de sciences biologiques, Université de Montréal, C.P. 6128, succursale Centre-ville,  
Montréal, Québec, Canada H3C 3J7

<sup>2</sup> Corresponding address: Daniel Borcard, Département de sciences biologiques, Université de  
Montréal, C.P. 6128, succursale Centre-ville, Montréal, Québec, Canada H3C 3J7. E-mail:  
Daniel.Borcard@umontreal.ca

Manuscript received 22 September 2011; revised 1 February 2012. Corresponding Editor: Brian  
Inouye.

1 *Abstract.* The Mantel correlogram is an elegant way to compute a correlogram for multivariate data.  
2 However, recent papers raised concerns about the power of the Mantel test itself. Hence the question: Is  
3 the Mantel correlogram powerful enough to be useful? To explore this issue, we compared the  
4 performances of the Mantel correlogram to those of other methods, using numerical simulations based  
5 on random, normally distributed data. For a single response variable, we compared it to the Moran and  
6 Geary correlograms. Type I error rates of the three methods were correct. Power of the Mantel  
7 correlogram was nearly as high as that of the univariate methods. For the multivariate case, the test of  
8 the multivariate variogram developed in the context of multiscale ordination is in fact a Mantel test, so  
9 that the power of the two methods is the same by definition. We devised an alternative permutation test  
10 based on the variance, which yielded similar results. Overall, the power of the Mantel test was high, the  
11 method successfully detecting spatial correlation at rates similar to the permutation test of the variance  
12 statistic in multivariate variograms. We conclude that the Mantel correlogram deserves its place in the  
13 ecologist's toolbox.

14 *Key words: Mantel; Geary; Moran; correlogram; multiscale ordination; spatial correlation;*  
15 *autocorrelation; multivariate variogram; simulation study*

16

17

18

## INTRODUCTION

A traditional way of measuring spatial correlation in a univariate quantitative variable is to resort either to Moran's  $I$  (1950) or Geary's  $c$  (1954) spatial correlation statistics (Cliff and Ord 1981). These structure functions, each in its manner, measure the resemblance of pairs of values located within predefined distance classes. Practically, one computes a matrix of geographical distances among sampling sites and converts these distances into a vector of classes  $\mathbf{d}$ . The statistic is then computed in turn within each distance class  $d$ .

Moran's  $I$  formula bears a close resemblance to the Pearson correlation coefficient:

$$I(d) = \frac{\frac{1}{W} \sum_{h=1}^n \sum_{i=1}^n w_{hi} (y_h - \bar{y})(y_i - \bar{y})}{\frac{1}{n} \sum_{i=1}^n (y_i - \bar{y})^2} \text{ for } h \neq i \quad (1)$$

whereas Geary's  $c$  is akin to a squared distance measure:

$$c(d) = \frac{\frac{1}{2W} \sum_{h=1}^n \sum_{i=1}^n w_{hi} (y_h - y_i)^2}{\frac{1}{n-1} \sum_{i=1}^n (y_i - \bar{y})^2} \text{ for } h \neq i \quad (2)$$

$y_h$  and  $y_i$  are the values of variable  $y$  at pairs of sites  $h$  and  $i$ ;  $\bar{y}$  is the mean of variable  $y$  over all sites.

Both formulas show the computation of the index value for a class of inter-site distance  $d$ .  $W$  is the number of pairs of points in the distance class considered. The weights  $w_{hi}$  have value 1 for pairs of sites belonging to distance class  $d$ , and 0 otherwise.

The expected value of Moran's  $I$  for no spatial correlation is  $E(I) = -1/(n-1)$ . For Geary's  $c$ , the expected value is 1.

For multivariate data, Sokal (1986) and Oden and Sokal (1986) proposed to construct a spatial correlation function on the basis of the Mantel (1967) test of matrix correlation. The principle goes as

1 follows: one computes an appropriate resemblance matrix  $\mathbf{X}$  among sites on the basis of the ecological  
 2 (or other) data, and for each geographical distance class this matrix is compared using the standardized  
 3 Mantel statistic  $r_M$  to a model matrix  $\mathbf{Y}$  where pairs of sites belonging to the same distance class  
 4 receive value 1 and the other pairs receive value 0. The resemblance matrix subjected to testing can be  
 5 computed in any way appropriate to the application field, e.g. using an ecological or genetic distance  
 6 function. It does not have to be computed using the Euclidean distance formula. The standardized  
 7 Mantel statistic has the same formula as the Pearson correlation coefficient, but computed between the  
 8 values in distance or similarity matrices  $\mathbf{X}$  and  $\mathbf{Y}$  (the lower- [or upper-]triangular values only are used  
 9 in the common case of a symmetric matrix  $\mathbf{X}$ ; the diagonal values are trivial and not used in the  
 10 calculation):

$$11 \quad r_M = \frac{1}{d-1} \sum_{i=2}^n \sum_{j=1}^{i-1} \left( \frac{x_{ij} - \bar{x}}{s_x} \right) \left( \frac{y_{ij} - \bar{y}}{s_y} \right) \quad (3)$$

12 where  $\bar{x}$  and  $\bar{y}$  are the means of the values in each of the lower-triangular distance matrices  $\mathbf{X}$  and  $\mathbf{Y}$   
 13 and  $s_x$  and  $s_y$  are their standard deviations.  $d = [n(n-1)/2]$  is the number of distances in the lower  
 14 triangular part of each distance matrix. The expected value for no correlation is 0.

15 Note that despite the mathematical relationship between Pearson's  $r$  and Mantel's  $r_M$ , one should  
 16 not try to interpret the actual  $r_M$  values as if they were equivalent to correlation coefficients among  
 17 variables. Dutilleul *et al.* (2000) showed that for pairs of simulated variables with normal distributions,  
 18 the  $r_M$  statistic computed between Euclidean distance matrices derived from these variables was always  
 19 much smaller than the theoretical correlation between the simulated variables. They explained this  
 20 phenomenon by showing, theoretically and empirically, that  $r_M$  calculated between matrices of squared  
 21 Euclidean distances computed from these variables is equal to the squared Pearson's  $r$  between the  
 22 original variables; the demonstration holds for the bivariate normal case only. When interpreting a  
 23 Mantel correlogram, one should not look for high values of the statistic, but for the shape drawn by the

1 significant  $r_M$  values. The sign of the coefficient is important, however. A positive (and significant)  $r_M$   
2 indicates that for the given distance class, the multivariate similarity among sites is higher than  
3 expected by chance (i.e., the mean within-class similarity is higher than the mean among-class  
4 similarity). The reverse is true for negative  $r_M$  values: communities that are at the corresponding  
5 distance are more different than expected by chance. This interpretation is valid for  $r_M$  statistics  
6 computed between an ecological *similarity* matrix and distance class model matrices with value 1 for  
7 pairs of sites belonging to distance class  $d$ , and 0 otherwise. If the calculation is based on an ecological  
8 *distance* matrix, the sign of  $r_M$  has to be changed for the correlogram to display positive values in cases  
9 of positive spatial correlation. This change of sign is done automatically in function `mantel.correlog()`  
10 in R.

11 Parametric tests exist for Moran's  $I$  and Geary's  $c$  coefficients. The Mantel statistic is generally  
12 tested by permutations, although a normal approximation exists for large data sets. Furthermore,  
13 Appendix A shows that a permutational test of Mantel's  $r_M$  on a matrix of squared Euclidean distances  
14 computed from univariate data and a distance model matrix is equivalent to a permutational test of  
15 Geary's  $c$  computed on the same data. It is also equivalent to a permutation test of the univariate semi-  
16 variance statistic used in variograms.

17 A correlogram is a graph representing the spatial correlation values against the distance classes.  
18 Correlograms can be drawn on the basis of any of the three functions described above. Depending on  
19 the way that the distance classes have been constructed, a correlogram can be all-directional (when the  
20 variable of interest is isotropic) or directional (when the geographic distances have been calculated in a  
21 geographic direction determined by the user). Interpretation of correlograms, which can be tricky, is  
22 explained by Legendre and Fortin (1989) and Legendre and Legendre (1998, 2012, Chapter 13).

23 Apart from the Mantel correlogram, techniques to assess multivariate spatial structures by means  
24 of structure functions are scarce. One interesting approach is multiscale ordination (MSO; Wagner

1 2003, 2004). MSO allows a spatial partitioning of community variation among distance classes and a  
2 representation and testing of the community spatial structure in the form of a multivariate variogram,  
3 which is simply the sum of the univariate variograms (Wagner 2003; Legendre and Legendre 2012,  
4 Chapter 13).

5 The Mantel correlogram is an elegant idea, but recent papers have raised concerns about the  
6 power of the Mantel test of matrix correlation itself (Legendre et al. 2005, Legendre et al. 2008,  
7 Legendre and Fortin 2010). Therefore, it is appropriate to investigate whether the Mantel correlogram  
8 has sufficient power to be useful to ecologists. This is why we set out to compare it to the performance  
9 of Moran's  $I$  and Geary's  $c$  in the univariate context (the use of a Mantel correlogram in the univariate  
10 context was needed only for comparison purposes), and to the multivariate variogram in the  
11 multivariate context, by using simulated data. We chose to base our simulations on random normal data  
12 compared using the Euclidean distance to provide a comparison of the Mantel correlogram with Moran  
13 and Geary correlograms in situations where the latter two are expected to perform at their best. The  
14 question addressed in this paper is thus: how does the Mantel correlogram compare to the other  
15 methods in terms of type I error rate and power in the case of random normal data and Euclidean  
16 distance matrices?

## 17 MATERIALS AND METHODS

### 18 *Data Simulations*

19 The shape of our simulated data surface was a square ( $100 \times 100$  pixel) grid populated by values  
20 drawn at random from the normal distribution, which were autocorrelated as described below.

21 A necessary preliminary step to a power study is to verify that the method under investigation has  
22 a correct rate of type I error. To that effect, we populated our grid with values drawn at random from a  
23 normal distribution with mean  $\mu = 0$  and standard deviation  $\sigma = 1$ . These values were generated using  
24 function `rnorm()` of the R language BASE package.

1           The random autocorrelated data were generated using an R function derived from the FORTRAN  
2 program SimSSD (Legendre et al. 2002, 2004, 2005). SimSSD is available as a Supplement to  
3 Legendre et al. (2005) (Ecological Archives # M075-017-S1).

4           Data generation was done as follows: random normal data were modified using conditional  
5 sequential Gaussian simulations based on a variogram, which produced data with isotropic spatial  
6 correlation. The spherical variogram model was controlled by values of the range parameter; values for  
7 this parameter were provided for each simulation run. In all simulations, the nugget variance was set to  
8 0. The range parameter of the variogram was set using the following values: 5, 10, 20, 50 and 100, i.e.,  
9 spatial correlation was made to vary among simulated data set from below the smallest range  
10 theoretically perceptible by the sampling design up to the point of encompassing the whole data set  
11 (100 units). The data were actually generated on a simulation grid larger than  $100 \times 100$  pixel to avoid  
12 border effects.

13           The sampling design was a regular  $10 \times 10$  grid overlaid on the  $100 \times 100$  grid; every tenth point  
14 was sampled in both directions. This resolution allowed us to sample data that were autocorrelated at  
15 scales starting well below the sampling interval (i.e., spatial correlation that should not be detected by  
16 analysis of the sample of observations), and ranging up to the broadest possible scale. Every simulation  
17 run consisted of 1000 independent replicate data with the same characteristics.

18           The multivariate data for the Mantel correlograms and multivariate variograms were made of two  
19 blocks of five independent variables. The first five variables were autocorrelated over the ranges  
20 defined above (except for the type I error rate simulations) whereas the last five variables were random  
21 normal variates with mean  $\mu = 0$  and standard deviation  $\sigma = 2$ . We added this amount of noise to the  
22 data to prevent the spatial correlation from being too strong and thus detectable with 100% power by  
23 all methods under investigation. All data sets, including the random normal variates, were generated

1 prior to their use in the test runs, so that the exact same data could be submitted to the different  
2 statistical methods.

### 3 *Computation and tests of spatial statistics*

4 All correlograms were computed for the same distance classes to make them comparable. The  
5 class interval, measured in units on the simulated surface, was equal to 8.6 pixels.

6 The line of reasoning developed in Appendix A led us to run two sets of Mantel tests for  
7 univariate data: one set where a matrix of Euclidean distances computed on the simulated data was  
8 used and another set where the distances were squared.

9 A set of computations consisted of the following steps: 1) Sample the next vector available  
10 among the 1000 data sets. 2) Compute the statistics (Moran's  $I$ , Geary's  $c$ , multivariate variogram and  
11 Mantel's  $r_M$ ) for the seven first distance classes defined above. 3) Test the coefficients. Variances in the  
12 variograms and correlations  $r_M$  in Mantel correlograms and in multivariate variograms were tested by  
13 999 permutations; the other two tests are parametric. 4) Write the values of the statistics and the  
14 associated probabilities to an output file. 5). Repeat steps 1 to 4 for the 1000 data sets. 6) Compute  
15 summary statistics: mean value of the statistics, rejection rates of  $H_0$  at  $\alpha = 0.05$  and their confidence  
16 intervals.

17 The Mantel correlogram was computed using the function `mantel.correlog()` of the VEGAN  
18 R package. Moran's  $I$  and Geary's  $c$  correlograms were obtained using a modified version of function  
19 `correlog()` of the package PGIRMESS. The multivariate variogram was obtained through a modified  
20 version of function `mso()` (VEGAN package). `correlog()` computes one-tailed tests in the direction  
21 of positive spatial correlation. `mso()` computes two-tailed tests. Therefore, we modified  
22 `correlog()` and `mso()` to allow the tests to be one-tailed in the direction of the sign of  $r_M$ , i.e., the  
23 same test as the default in `mantel.correlog()`. This may seem an overly liberal decision, but we

1 justify it by the fact that when computing a correlogram one often has one-tailed hypotheses of positive  
2 spatial correlation for the first few distance classes. For these classes, a two-tailed test would be too  
3 conservative. Conversely, many spatial structures induce *negative* spatial correlation at larger distances,  
4 which are detected by one-tailed tests in the direction of the sign (which detect negative as well as  
5 positive correlations). A side effect of this decision is that in the simulations for type I error, one  
6 expects the overall rejection rate to be twice the  $\alpha$  significance level. For instance, for  $\alpha = 0.05$ , a test is  
7 expected to reject  $H_0$  in 5% of the runs in each tail, i.e., in 10% of the runs.

8 In its current version, the variogram test provided by function `msO()` is in fact the exact same  
9 Mantel test as performed by the function `mantel.correlog()`. Consequently, we had no  
10 independent test to which we could compare the performance of the Mantel correlogram. This led us to  
11 devise another test based on a different statistic. The Mantel test is based on the correlation between a  
12 matrix of (ecological) distances among sites and a binary matrix representing the membership of sites  
13 to the distance class being tested. In our new test, the variogram statistic (variance) corresponding to  
14 each distance class is tested by random permutations of the data rows followed by recomputation of the  
15 statistic. The test is one-tailed in the direction of the sign of the difference between the statistic and the  
16 total variance of the data, which is the mathematical expectation of the variogram statistic. The test is  
17 run as follows for a given distance class  $k$ : (1) compute the variance of class  $k$ ,  $s^2_k$  and the total variance  
18  $s^2_{tot}$  of the data; (2) note the sign of the statistic obtained by subtracting  $s^2_{tot}$  from  $s^2_k$ ; (2) permute the  
19 rows of the data table; (3) recompute the variance of the permuted data in class  $k$ ,  $s^2_{k,perm}$ ; (4) compare  
20  $s^2_{k,perm}$  to the distribution of  $s^2_k$  obtained by permutation, in the tail given by the sign noted at step 2.

21 Another issue is the correction for multiple testing often applied to correlograms. Repeated  
22 testing on a given data set increases the overall risk of type I error. An overview of the most often-used  
23 correction procedures can be found in Legendre and Legendre (1998, p.18). In the simulations, we  
24 opted for no correction in order to obtain and compare the raw probabilities in each distance class.

## RESULTS

### *Type I error rate*

Fig. 1 shows the results of the type I error simulations for completely random, uncorrelated data in the univariate case. Fig. 4a and 4b provide the same type of results in the multivariate case; the Mantel correlogram and variance-based test of the multivariate variogram had very similar results. As can be seen, all methods investigated are valid for all distance classes. A statistical test is valid when the rejection rate of  $H_0$  is not larger than the significance level  $\alpha$ , for any value of  $\alpha$ , when  $H_0$  is true (Edgington 1995). Validity has been verified for other  $\alpha$  levels as well (not illustrated). Type I error rates were always correct (i.e., close to the expected value of 10% in our simulations). All statistics showed values close to their expectation at all distance classes. The results are the same for data that were autocorrelated with a range of 5 units, i.e., half the sampling interval (not illustrated). This is normal, since the spatial correlation range was below the detection limit of the sampling design. In spite of the development presented in Appendix A, there are slight differences between the Mantel and Geary results, due to the fact that the tests of Geary correlograms were parametric.

### *Power*

The results reported below always refer to the Mantel correlogram computed on squared Euclidean distances because this version of the  $r_M$  statistic is mathematically closer to Geary's  $c$ , as demonstrated in Appendix A. Small but nonsignificant differences between results based on Euclidean distances and squared Euclidean distances are due to the randomness of the permutations.

### **Univariate data**

Overall and as expected, the power of the three methods increased with the range of the spatial correlation in the data. The power of the three methods was approximately the same, with a slight advantage to Moran and Geary (for which our simulations used parametric tests, compared to permutational tests for the Mantel correlogram).

1 Fig. 2a to 2d show the results of the simulations for data autocorrelated with a range of 10 units,  
2 i.e., corresponding to the sampling interval. Interestingly, over the 1000 independent runs, the three  
3 coefficients detected spatial correlation at the first distance class enough times to change the mean  
4 values of the statistics by a very slight amount (Fig. 2a, 2b and 2c, where the scale of the ordinate was  
5 chosen to show tiny differences), but the power of these methods did not reach over the level of type I  
6 error (Fig. 2d). This means that even in the cases where these three methods reacted to short-range  
7 spatial correlation with a higher coefficient value, these values were not more often significant than  
8 with uncorrelated data. This is due to the fact that, in order to be consistently detected by the test,  
9 spatial correlation has to be larger than the class interval.

10 Increasing the range of spatial correlation increased the rates of detection at short range (first  
11 distance classes) and at progressively larger ranges. With a spatial correlation range of 20 units (Fig. 2e  
12 to 2h), the powers of the Moran and Geary coefficients for the first distance class were near 0.7.  
13 Mantel's  $r$  was slightly less powerful, at 0.6 (Fig. 2h). The power of the three coefficients dropped to  
14 around or below 0.20 for the second distance class, and remained slightly above the type I error level  
15 for the other five classes.

16 With a spatial correlation range of 50 units (Fig. 3a to 3d), the power of the three statistics was  
17 high for the first two distance classes (close to 0.90; Fig. 3d). Spatial correlation was detected in  
18 approximately 40% of the cases in the third distance class. The rate dropped to around 0.3 in class 4,  
19 but increased slightly in the next classes, which displayed negative spatial correlation. When the spatial  
20 correlation range reached the extent of the simulated surface (range = 100), the results (Fig. 3e to 3h)  
21 resembled those obtained for a range of 50 units, but with the smallest rates of rejection in distance  
22 class 5 (Fig. 3h), and negative spatial correlation detected in around 40 to 55% of the cases in classes 6  
23 and 7.

1           The conclusion from this part of the study is that the three methods performed in a very  
2 comparable manner, the Mantel correlogram being only slightly less powerful for univariate normal  
3 data than its classical univariate parametric counterparts.

#### 4 **Multivariate data**

5           For reasons explained in the Materials and Methods section, the type I error and rejection rates of  
6 the Mantel correlogram and of the multivariate variogram in the form implemented in the version of  
7 `msO()` currently distributed with `VEGAN` on `CRAN` were identical to within a few random  
8 permutations. This equivalence holds because the test used in `msO()` is a Mantel test based on  
9 Euclidean distances. A Mantel test based on a matrix of non-Euclidean (e.g., Bray-Curtis) ecological  
10 distances would of course not be equivalent to a multivariate variogram computed on raw data.  
11 Normality of distributions, however, plays no role in this equivalence because the tests are  
12 permutational. Furthermore, the power of our new test of the multivariate variogram, based on  
13 variances, was undistinguishable from that of the Mantel correlogram for all spatial correlation ranges  
14 and distance classes. Therefore, only the results for the Mantel correlogram are presented.

15           Fig. 4c to 4j show the distributions of  $r_M$  statistics and rejection rates of the Mantel correlogram  
16 for four ranges of spatial correlation. Overall, the properties observed here are the same as the ones  
17 displayed by the univariate Mantel correlogram. As soon as the range of spatial correlation exceeds the  
18 sampling distance, the power of the Mantel correlogram increases, first in the first distance class, and  
19 then in the following ones when the range increases. Note the power to detect negative spatial  
20 correlation, lying at about 40% in class 7 of ranges 50 (Fig. 4h) and 100 (Fig. 4j).

21           The conclusion of this part of the study is that the Mantel test in multivariate correlogram  
22 analysis had the same type I error rate and power as the permutation test of the new variogram statistic.  
23 Furthermore, it displayed a power comparable to those of the univariate methods for normally  
24 distributed data with similar ranges of spatial correlation.

## CONCLUSION

1  
2 All results reported above are dependent on the relationship between the range of the spatial  
3 patterns or correlation and the grain, i.e. the width of the distance classes chosen in the analyses. In this  
4 study based on a square grid with intersite distances equal to 10, we chose a class interval (8.6) that  
5 allowed the first class to encompass sites that were related horizontally, vertically and diagonally, since  
6 the interval [8.6, 17.4) contains the diagonal distance  $\sqrt{200} = 14.14$  (remember, the points are 10 units  
7 apart horizontally and vertically). The second class contained the second neighbors, and so on. It was  
8 important to choose a class interval that was both compatible with the grain of the study and the aim of  
9 the analysis. Since this condition was fulfilled, our study showed that, despite the recent concerns about  
10 the power of the Mantel test, which is at the core of the Mantel correlogram, this method provided a  
11 reliable assessment of the presence of spatial correlation in univariate as well as multivariate data. The  
12 concerns expressed in recent literature about the Mantel test are related to situations where alternative  
13 methods are available to address the scientific hypotheses more appropriately and to test them with  
14 more power. When looking for a multivariate assessment of spatial correlation, however, few  
15 alternatives exist to compute correlograms, which test hypotheses formulated in terms of distances.  
16 Therefore, our simulation results can be seen from two viewpoints: (1) comparisons between the  
17 Mantel correlogram and other approaches using data compatible with all methods showed that the  
18 Mantel correlogram behaves well; (2) in its own field, in the absence of equivalent methods, the Mantel  
19 correlogram has an acceptable power. For univariate data, the test of significance in the Mantel  
20 correlogram based on a Euclidean distance matrix is equivalent to the test of Geary's  $c$ , which is also  
21 the method to test autocorrelation values in univariate variograms. For multivariate data, the test used  
22 in Mantel correlograms has the exact same power as the only alternative method, which is the test of  
23 the multivariate variogram.

1           To be able to compare the methods with data compatible with all of them, we ran our simulations  
2 on random normal variables projected on sites that were compared using the Euclidean distance, this  
3 class including distances that can be obtained by transforming the data followed by calculation of the  
4 Euclidean distance formula; see Legendre and Gallagher (2001) and Legendre and Legendre (2012,  
5 Section 7.7). However, the real value of the Mantel correlogram is for studies where the response data  
6 are modeled by a distance function that is not Euclidean. Among these are most of the coefficients  
7 developed for the analysis of species presence-absence data, for example the coefficients of Jaccard  
8 and Sørensen, as well as quantitative distances for species abundance data, such as the asymmetric  
9 Gower coefficient, the geodesic metric, the Whittaker, Canberra, Clark, percentage difference, and  
10 mean character difference modified for species data. Distance coefficients intended for data other than  
11 species are, for example, the symmetric Gower, Estabrook-Rogers, and the generalized Mahalanobis  
12 distance for groups of observations. See Legendre and Legendre (2012) for descriptions of these  
13 coefficients, which can all be used in Mantel correlogram studies and in none of the other methods.  
14 Further simulations are needed to assess the behavior of the Mantel correlogram with non-normal data  
15 and non-Euclidean distances.

16

#### ACKNOWLEDGMENTS

17           We are grateful to Alena Motorina for help at the beginning of the simulation process, and to two  
18 anonymous reviewers for very insightful comments on the manuscript. This work was supported by  
19 Natural Sciences and Engineering Research Council of Canada (NSERC) grant no. 7738 to P.  
20 Legendre.

21

## LITERATURE CITED

- 1
- 2 Cliff, A.D., and J. K. Ord. 1981. Spatial processes. Pion, London.
- 3 Dutilleul, P., J. D. Stockwell, D. Frigon and P. Legendre. 2000. The Mantel test versus Pearson's  
4 correlation analysis: assessment of the differences for biological and environmental studies.  
5 Journal of Agricultural, Biological and Environmental Statistics 5: 131-150.
- 6 Edgington, E. S. 1995. Randomization tests. 3rd edition. Marcel Dekker, Inc., New York.
- 7 Geary, R. C. 1954. The contiguity ratio and statistical mapping. The Incorporated Statistician 5: 115-  
8 145.
- 9 Giraudoux, P. 2010. pgirmess: Data analysis in ecology. R package version 1.4.7. [http://CRAN.R-](http://CRAN.R-project.org/package=pgirmess)  
10 [project.org/package=pgirmess](http://CRAN.R-project.org/package=pgirmess).
- 11 Legendre, P., D. Borcard, and P. R. Peres-Neto. 2005. Analyzing beta diversity: partitioning the spatial  
12 variation of community composition data. Ecological Monographs 75: 435-450.
- 13 Legendre, P., D. Borcard, and P. R. Peres-Neto. 2008. Analyzing or explaining beta diversity:  
14 Comment. Ecology 89: 3238-3244.
- 15 Legendre, P., M. R. T. Dale, M.-J. Fortin, J. Gurevitch, M. Hohn, and D. Myers. 2002. The  
16 consequences of spatial structure for the design and analysis of ecological field surveys.  
17 Ecography 25: 601-615.
- 18 Legendre, P., M. R. T. Dale, M.-J. Fortin, P. Casgrain, and J. Gurevitch. 2004. Effects of spatial  
19 structures on the results of field experiments. Ecology 85: 3202-3214.
- 20 Legendre, P. and M.-J. Fortin. 1989. Spatial pattern and ecological analysis. Vegetatio 80: 107-138.

- 1 Legendre, P., and M.-J. Fortin. 2010. Comparison of the Mantel test and alternative approaches for  
2 detecting complex multivariate relationships in the spatial analysis of genetic data. *Molecular*  
3 *Ecology Resources* 10: 831-844.
- 4 Legendre, P., and L. Legendre. 1998. *Numerical ecology*, 2nd English edition. Elsevier Science BV,  
5 Amsterdam.
- 6 Legendre, P., and L. Legendre. 2012. *Numerical ecology*, 3rd English edition. Elsevier Science BV,  
7 Amsterdam.
- 8 Mantel, N. 1967. The detection of disease clustering and a generalized regression approach. *Cancer*  
9 *Research* 27: 209-220.
- 10 Moran, P. A. P. 1950. Notes on continuous stochastic phenomena. *Biometrika* 37: 17-23.
- 11 Oden, N. L., and R. R. Sokal 1986. Directional autocorrelation: an extension of spatial correlograms to  
12 two dimensions. *Systematic Zoology* 35: 608-617.
- 13 Oksanen, J., G. Blanchet, R. Kindt, P. Legendre, B. O'Hara, G. L. Simpson, P. Solymos, M. H. H.  
14 Stevens, and H. Wagner. 2011. *vegan: Community Ecology Package*. R package version 1.17-11.  
15 <http://cran.r-project.org/package=vegan>.
- 16 Sokal, R. R. 1986. Spatial data analysis and historical processes. Pp. 29-43 in: E. Diday et al. (eds.)  
17 *Data analysis and informatics, IV*. North-Holland, Amsterdam.
- 18 Wagner, H. H. 2003. Spatial covariance in plant communities: integrating ordination, variogram  
19 modeling, and variance testing. *Ecology* 84: 1045–1057.
- 20 Wagner, H. H. 2004. Direct multi-scale ordination with canonical correspondence analysis. *Ecology*  
21 85: 342–351.
- 22

## 1 FIGURE CAPTIONS


2 Fig. 1 Values of the statistics (a, b, c), type I error rates (d) and 95% confidence intervals for seven  
 3 distance classes of Moran (circles), Geary (squares) and Mantel (diamonds) correlograms run on  
 4 random normal data (1000 independent runs). The Mantel correlograms were computed on squared  
 5 distances. Type I error rates are for a significance level  $\alpha = 0.05$ , one-tailed on the side of the sign. The  
 6 expected rejection rate under  $H_0$  was thus 0.10. Horizontal lines in the graphs indicate the expectation  
 7 of the statistic (a, b, c) or the type I error rate (d).

8 Fig. 2 Values of the statistics of Moran's  $I$  (circles; a, e), Geary's  $c$  (squares; b, f) and Mantel's  $r$   
 9 (diamonds; c, g) correlograms and rejection rates of  $H_0$  (d, h) for normal data autocorrelated with a  
 10 range of 10 units (a, b, c, d) and 20 units (d, e, f, g) on the simulated surface and sampled every tenth  
 11 unit in both directions. Rejection rates are for a significance level  $\alpha = 0.05$ , one-tailed on the side of the  
 12 sign. The expected rejection rate under  $H_0$  was thus 0.10. Horizontal lines in the graphs indicate the  
 13 expectation of the statistic (a, b, c, e, f, g) or the expected rejection rate under  $H_0$  (d, h). 95%  
 14 confidence intervals are omitted when they are smaller than the symbols.


15 Fig. 3 Values of the statistics of Moran's  $I$  (circles; a, e), Geary's  $c$  (squares; b, f) and Mantel's  $r$   
 16 (diamonds; c, g) correlograms and rejection rates of  $H_0$  (d, h) for normal data autocorrelated with a  
 17 range of 50 units (a, b, c, d) and 100 units (d, e, f, g) on the simulated surface and sampled every tenth  
 18 unit in both directions. For details, see caption of Fig. 2.

19 Fig. 4 Values of the statistics (a, c, e, g, i), type I error rates (b), power (which is the rejection rate of  $H_0$   
 20 when spatial correlation is present) (d, f, h, j) and 95% confidence intervals for seven distance classes  
 21 of multivariate Mantel correlograms run on random normal data (a, b) and spatially correlated data (c  
 22 to j) in each of 1000 independent runs. Type I error and rejection rates: see caption of Fig. 2.


**Fig. 1**





**Fig. 2**



**Fig. 3**



**Fig. 4****Mantel's  $r$** **Rejection rates,  $r_M$** 

## APPENDIX A

*Ecological Archives* Exxx-xxx-A1

### MATHEMATICAL LINK BETWEEN THE PERMUTATIONAL TESTS OF GEARY'S $C$ AND MANTEL'S $Z_M$

This Appendix presents a demonstration of the equivalence between the permutational tests of the semi-variance, Geary's  $c$ , and Mantel's  $z_M$  statistics.

## MANTEL'S Z AND GEARY'S C ARE RELATED

We demonstrate here that in the form used in the Mantel correlogram, the original Mantel statistic  $z_M$  is closely related to Geary's  $c$  and should produce the exact same permutational results when the Mantel correlogram is computed on squared distances. The demonstration concerns  $z_M$  instead of the widely used standardized coefficient  $r_M$ , but in practical terms this is not important since both forms also produce the exact same permutational test results. The comparison is made for the analysis of a single variable and the distance used to compute the Mantel correlogram statistic is the Euclidean distance.

1. Consider a matrix  $\mathbf{D}$  of Euclidean distances computed on the variable of interest. In the Mantel correlogram, for a given class of geographical distance  $d$ ,  $z_M$  is the scalar product of matrix  $\mathbf{D}$  and a binary model matrix built as explained in the Introduction, after transforming (stringing out) both matrices into vectors. Only the upper (or lower) triangular portion of each matrix need be used in the calculation since a Euclidean distance matrix is symmetric. Therefore,  $z_M$  is simply the sum of the distances of  $\mathbf{D}$  that belong to distance class  $d$ .

2. The semi-variance (Legendre and Legendre 1998 eq. 13.10) is the numerator of Geary's  $c$ :

$$\gamma(d) = \frac{1}{2W} \sum_{h=1}^{n-1} \sum_{i=h+1}^n w_{hi} (y_h - y_i)^2 \quad (4)$$

- Therefore, a permutational test of Geary's  $c$  is equivalent to a permutational test of  $\gamma$  since the two statistics differ only by the denominator of  $c$ , which is a constant.
- $w_{ih}$  is used to select the pairs belonging to class  $d$ ; it plays a role equivalent to the binary model matrix in the Mantel correlogram test.

- $W$  is the numbers of pairs in distance class  $d$ ; since this number is constant across the permutations, one can eliminate it without changing the result of the test.
- The computation of Mantel's  $z_M$  differs from  $\gamma$  by a single point: to obtain  $z_M$ , one sums the Euclidean distances between pairs of objects, whereas to obtain  $\gamma$  one sums *squared* Euclidean distances.

**Conclusion:** one can obtain a test equivalent to a *permutational* test of the semi-variance statistic or Geary's  $c$  by squaring the distances in  $\mathbf{D}$  before submitting them to the Mantel test.